

TAME COVERS AND COHOMOLOGY OF RELATIVE CURVES OVER COMPLETE DISCRETE VALUATION RINGS WITH APPLICATIONS TO THE BRAUER GROUP

ERIC BRUSSEL AND EDUARDO TENGAN

ABSTRACT. We prove the existence of noncrossed product division algebras and indecomposable division algebras of unequal period and index over the function field of any p -adic curve, extending the results and methods of [10].

1. INTRODUCTION

We study the cohomology and the Brauer group of a field F that is finitely generated and of transcendence degree one over the p -adic field \mathbb{Q}_p . Such a field is always the function field of a regular (projective, flat) relative curve X/\mathbb{Z}_p . In [10] it was shown that if F admits a *smooth* model X/\mathbb{Z}_p then there exist noncrossed product F -division algebras, and indecomposable F -division algebras of unequal prime-power period and index. These were constructed from objects defined over the generic point (p) of the closed fiber $X_0 = X \otimes_{\mathbb{Z}_p} \mathbb{F}_p$ using a homomorphism $\lambda : \text{Br}(F_p)' \rightarrow \text{Br}(F)$ that splits the restriction map $\text{res} : \text{Br}(F) \rightarrow \text{Br}(F_p)$, where the subscript p denotes completion with respect to the valuation defined by (p) , and “ $'$ ” denotes “prime-to- p part”. The fields F that have a smooth X/\mathbb{Z}_p include fields such as $\mathbb{Q}_p(t)$ but do not include the function fields of *all* p -adic curves.

In this paper we generalize the machinery and results of [10] to arbitrary p -adic curves. We prove that if F is the function field of a p -adic curve then there exist noncrossed product F -division algebras, and indecomposable F -division algebras of unequal prime-power period and index. The machinery we develop here is used in [11] to prove that every F -division algebra of prime period ℓ and index ℓ^2 decomposes into two cyclic F -tensor factors, hence is a crossed product, generalizing Suresh’s result [36], which assumes roots of unity. In the terminology of [5, Sections 3,4], this shows the \mathbb{Z}/ℓ -length in $H^2(F, \mu_\ell)$ equals the ℓ -Brauer dimension, which is two by a theorem of Saltman ([33, Theorem 3.4]). In general our work is motivated by the work of Saltman over these fields in [33] and [35] (see also [9]).

We summarize the technical results. Let R be a complete discrete valuation ring with fraction field K , and let F be a finitely generated field extension of K of transcendence degree one. Let X/R be a regular (projective, flat) model for F whose closed fiber X_0 has normal crossings on X . Let $C = X_{0,\text{red}}$, let $\{C_i\}$ denote the set of irreducible components of C , let \mathcal{S} be the set of singular points of C , and set $F_C = \prod_i F_{C_i}$, the product of the completions with respect to the valuations

1991 *Mathematics Subject Classification.* 11G20, 11R58, 12G05, 16K20, 16K50.

on F defined by the C_i . We construct for any prime-to-char($\kappa(C)$) number n , any integer r , and any $q \geq 0$, a homomorphism

$$\lambda : H^q(O_{C,S}, \mathbb{Z}/n(r)) \rightarrow H^q(O_{X,S}, \mathbb{Z}/n(r)) \rightarrow H^q(F, \mathbb{Z}/n(r))$$

that splits the restriction map $H^q(O_{X,S}, \mathbb{Z}/n(r)) \rightarrow H^q(O_{C,S}, \mathbb{Z}/n(r))$. We use λ to construct another map λ that splits the subgroup of the image of $H^q(F, \mathbb{Z}/n(r)) \rightarrow H^q(F_C, \mathbb{Z}/n(r))$ consisting of tuples of classes that are unramified at \mathcal{S} and glue along \mathcal{S} . When $q = 2$, $\mathbb{Z}/n(r) = \mu_n$, and $R = \mathbb{Z}_p$, we show that λ preserves index. This allows us to construct indecomposable F -division algebras and noncrossed product F -division algebras as mentioned above, in the same manner as [10]. When the dual graph of the closed fiber X_0 has nontrivial topology, i.e., nonzero (first) Betti number, we construct cyclic covers of X that are (defined and) trivial at every point of X except the generic point of X . These arise as cyclic covers of the closed fiber X_0 that are trivial at every point, and transported to X via λ . When $R = \mathbb{Z}_p$ they are the completely split cyclic covers of Saito ([30]). We thank Colliot-Thélène for drawing our attention to these interesting specimens.

2. BACKGROUND AND CONVENTIONS

We use [26, Chapter 8,9], [23, Section 2], and [22, Chapter XIII] for many of the following definitions.

2.1. General Conventions. Let S be an excellent scheme, n a number that is invertible on S , and $\Lambda = (\mathbb{Z}/n)(i)$ the twisted étale sheaf. We write $H^q(S, \Lambda)$ for the étale cohomology group, and if Λ is understood (or doesn't matter) we write $H^q(S, r)$ instead of $H^q(S, \Lambda(r))$, and $H^q(S)$ in place of $H^q(S, 0)$. If $S = \text{Spec } A$ for a ring A then we write $H^q(A, r)$. If T is an integral scheme contained in S we write $\kappa(T)$ for its function field. If $T \rightarrow S$ is a morphism of schemes then the restriction $\text{res}_{T|S} : H^q(T) \rightarrow H^q(S)$ is defined, and we write $\beta_S = \text{res}_{T|S}(\beta)$. If $Z \subset S$ is a subscheme, we write Z_T for the preimage $Z \times_S T$.

2.2. Valuation Theory. If v is a valuation on a field F we write $\kappa(v)$ for the residue field of the valuation ring O_v , and F_v for the completion of F at v . If S is a connected normal scheme with function field F and v arises from a prime divisor D on S , we write v_D for v , $\kappa(D)$ for $\kappa(v)$, and F_D for F_v . If D is a sum of prime divisors D_i we write $F_D = \prod_i F_{D_i}$. Each $f \in F^*$ defines a divisor $\text{div}(f) = \sum v_D(f)D$, where the (finite) sum is over prime divisors on S .

Recall that if $F = (F, v)$ is a discretely valued field and $\alpha \in H^q(F, \Lambda)$ then α has a *residue* $\partial_v(\alpha)$ in $H^{q-1}(\kappa(v), \Lambda(-1))$. More generally, suppose T is a noetherian scheme, ξ is a generic point of T , and $\alpha \in H^q(T, \Lambda)$. Then for each discrete valuation v on the field $F = \kappa(\xi_{\text{red}})$ α has a residue

$$\partial_v(\alpha) \stackrel{\text{df}}{=} \partial_v(\alpha_F) \in H^{q-1}(\kappa(v), \Lambda(-1))$$

We say α is *unramified at v* if $\partial_v(\alpha) = 0$, *ramified at v* if $\partial_v(\alpha) \neq 0$, and *tamely ramified at v* if $\partial_v(\alpha)$ is contained in the prime-to-char($\kappa(v)$) part of $H^{q-1}(\kappa(v), \Lambda(-1))$. If α is unramified at v the *value* of α at v is the element

$$\alpha(v) = \text{res}_{F|F_v}(\alpha_F) \in H^q(\kappa(v), \Lambda) \leq H^q(F_v, \Lambda)$$

(see [16, 7.13, p.19]). Suppose $T \rightarrow S$ is a birational morphism of noetherian schemes (see [17, I.2.2.9]). The *ramification locus* of α on S_{red} is the sum of the

prime divisors on S_{red} that determine valuations at which α is ramified, over all generic points of S_{red} .

Let D be a divisor on a noetherian normal scheme S , set $U = S - D$, and for each generic point ξ of $\text{Supp } D$, let K_ξ denote the fraction field of the discrete valuation ring $\mathcal{O}_{S,\xi}$. We say a morphism $\rho : T \rightarrow S$ is *tamely ramified along D* if T is normal, $\rho_U : V = T \times_S U \rightarrow U$ is étale, and for each generic point ξ of $\text{Supp } D$, the étale K_ξ -algebra L defined by $\text{Spec } L = V \times_U \text{Spec } K_\xi$ is tamely ramified with respect to $\mathcal{O}_{S,\xi}$. Since L/K_ξ is étale it is a finite product of separable field extensions of K_ξ , and L is tamely ramified if each field extension is tamely ramified (with respect to $\mathcal{O}_{S,\xi}$) in the usual sense. If S is a noetherian scheme whose irreducible components are normal, we'll say a morphism $\rho : T \rightarrow S$ is tamely ramified along D if again $V \rightarrow U = S - D$ is étale, and the restriction ρ_{S_i} to each irreducible component S_i of S is tamely ramified along D_{S_i} . If $S = \text{Spec } A$ and $T = \text{Spec } B$, we will also say B is a *tamely ramified A -algebra*. We say a map $\rho : T \rightarrow S$ of noetherian schemes is a *cover* if it is finite, generically étale, and each connected component of T dominates a connected component of S .

2.3. Relative Curves. In this paper, R will be a complete discrete valuation ring with residue field k and fraction field K , F will be a field finitely generated of transcendence degree one over K , and X/R will be a regular 2-dimensional scheme X that is flat and projective over $\text{Spec } R$ and has function field $K(X) = F$. We call X/R a *regular relative curve*, write $X_0 = X \otimes_R k$ for the closed fiber, $C = X_{0,\text{red}}$ for the reduced subscheme underlying the closed fiber, and C_1, \dots, C_m for the irreducible components of C . We assume each C_i is regular, and at most two of them meet at any closed point of X , a situation that can always be obtained by blowing up using Lipman's embedded resolution theorem (see [26, 9.2.4]). For all closed points $z \in X$, we have $\dim \mathcal{O}_{X,z} = 2$ by [26, 8.3.4(c)], and since X is regular, $\mathcal{O}_{X,z}$ is factorial by Auslander-Buchsbaum's theorem.

We say an effective divisor D on a relative curve X/R is *horizontal* if each of its irreducible components maps surjectively (hence finitely) to $\text{Spec } R$, and *vertical* if its support is contained in the support of the closed fiber. If D is a reduced and irreducible horizontal divisor then it is flat over $\text{Spec } R$, since R is a discrete valuation ring. Every effective divisor on a relative curve X/R is a sum of horizontal and vertical divisors, and the horizontal prime divisors are exactly the closures of the closed points of the generic fiber ([26, 8.3.4(b)]). Since R is henselian, each irreducible horizontal divisor has a single closed point.

2.4. Distinguished Divisors. In general there will be many horizontal divisors on a relative curve X that restrict to a given divisor on C . In order to construct our lifts of covers and cohomology classes from C to X we select a single regular horizontal divisor for each closed point, as follows.

Proposition 2.5. *Assume the setup of (2.3). Then for each closed $z \in X$ there exists a regular irreducible horizontal divisor $D \subset X$ that intersects each irreducible component of C passing through z transversally at z .*

Proof. Transversal intersection with a single component is by [26, 8.3.35(g)] and its proof (see also [21, 21.9.12]). Thus if $z \in C_i \cap C_j$ ($i \neq j$) and t_i and t_j are local equations for C_i and C_j , then we have local equations f_i and f_j for effective

regular horizontal divisors such that $(f_i, t_i) = (f_j, t_j) = \mathfrak{m}_z \subset \mathcal{O}_{X,z}$. If $(f_j, t_i) = \mathfrak{m}_z$ or $(f_i, t_j) = \mathfrak{m}_z$ then a suitable D is defined locally by f_j or f_i . Otherwise $(f_i + f_j, t_i) = (f_i + f_j, t_j) = \mathfrak{m}_z$, and we define D locally by $f_i + f_j$. The rest of the proof proceeds as in [26, 3.3.35]. \square

We fix a set of these (prime) divisors, and let \mathcal{D} denote the set of supports of the semigroup they generate in $\text{Div } X$. We will say a divisor D is *distinguished* and write $D \in \mathcal{D}$ whenever D is reduced and supported in \mathcal{D} . Though \mathcal{D} is fixed in principle, we reserve the right to declare any divisor satisfying the definition to be a member of this set retroactively. Let $\mathcal{D}_{\mathcal{S}}$ denote the subset that *avoids* \mathcal{S} . Note that each $D \in \mathcal{D}$ is a disjoint union of its irreducible components, each of which meets each irreducible component of C transversally.

3. STRUCTURE OF TAME COVERS

Lemma 3.1 (Structure). *Assume the setup of (2.3). Suppose $\rho : Y \rightarrow (X, D)$ is a tamely ramified cover, where $D \in \mathcal{D}$. Then*

- a) *The structure map $\rho : Y \rightarrow X$ is flat.*
- b) *Y/R is a regular relative curve, $Y_{0,\text{red}} = C_Y$, each irreducible component of C_Y is regular, \mathcal{S}_Y is the set of singular closed points of C_Y , and exactly two irreducible components of C_Y meet at each point of \mathcal{S}_Y .*
- c) *The support of the irreducible components of D'_Y for $D' \in \mathcal{D}$ generate a set \mathcal{D}_Y of distinguished divisors on Y .*

Proof. Since $Y \rightarrow X$ is finite, $\dim(X) = \dim(Y) = 2$ by [19, 5.4.2], and $Y \rightarrow \text{Spec } R$ is projective as the composition of projective morphisms ([26, 3.3.32]). Let $y \in Y$ be a closed point and set $x = f(y)$, $A = \mathcal{O}_{X,x}$, $B' = \mathcal{O}_{Y,x}$, and $B = \mathcal{O}_{Y,y}$. Choose a geometric point over x that lifts to each point of Y lying over x , and use this in the following to define the strict henselizations with respect to the maximal ideals of these points.

Since the statements involving D are local and D is a disjoint union of its irreducible components we may assume D is irreducible. Let $C_i \subset C$ be a (regular) irreducible component going through x , and let $\{f, t\} \subset A$ be the regular system of parameters formed by local equations for the distinguished prime divisor passing through x , and for C_i , respectively. Then the strict henselization A^{sh} of A with respect to the maximal ideal of A is a two-dimensional regular local ring, faithfully flat over A , with regular system of parameters $\{f, t\}$ (see [21, 18.8]).

If $x \notin D$ then $B' \otimes_A A^{\text{sh}}$ is a finite étale A^{sh} -algebra by base change, since $\rho|_{X-D}$ is finite-étale. If $x \in D$ then $B' \otimes_A A^{\text{sh}}$ is a finite tamely ramified A^{sh} -algebra by [23, Lemma 2.2.8]. By [21, 18.8.10], B^{sh} is a factor of the direct product decomposition of $B' \otimes_A A^{\text{sh}}$, hence B^{sh} is a finite tamely ramified local A^{sh} -algebra, in particular it is a normal local ring, hence it is a normal domain. It follows that B^{sh} is the integral closure of A^{sh} in the field $\tilde{L} \stackrel{\text{df}}{=} \text{Frac } B^{\text{sh}}$. Since the tame fundamental group of the strictly henselian regular local ring A^{sh} is abelian ([22, XIII.5.3]) the field extension $\tilde{L}/\text{Frac } A^{\text{sh}}$ is Galois, and by Abhyankar's Lemma ([13, A.I.11], see also [23, Corollary 2.3.4])

$$B^{\text{sh}} = A^{\text{sh}}[T]/(T^e - f) \quad (\text{some } e \geq 1)$$

By [23, Lemma 1.8.6] B^{sh} is a regular (2-dimensional) local ring with system of parameters $\{\sqrt[e]{f}, t\}$. Since $B \rightarrow B^{\text{sh}}$ is faithfully flat and B^{sh} is regular, B is regular by flat descent ([19, 6.5.1] or [27, 23.7(i)]), and since B is the local ring of an arbitrary closed point, we conclude Y is regular. It follows that $\rho : Y \rightarrow X$ is flat by [27, 23.1], proving (a), and since Y is regular and $Y \rightarrow \text{Spec } R$ is flat and projective, Y/R is a regular relative curve.

We derive a system of parameters for B . The prime ideal $(\sqrt[e]{f}) \subset B^{\text{sh}}$ is the only one lying over $(f)A^{\text{sh}}$ since, for $\kappa(f) = \text{Frac } A^{\text{sh}}/(f)A^{\text{sh}}$, the ring $B^{\text{sh}} \otimes_{A^{\text{sh}}} \kappa(f) = \kappa(f)[T]/(T^e)$ of the fiber over $\text{Spec } \kappa(f)$ consists of a single prime ideal. The image $(\sqrt[e]{f})$ in $\text{Spec } B$ is therefore a unique prime $(g) \subset B$ lying over $(f) \subset A$, and $(\sqrt[e]{f})$ is the unique prime lying over (g) . Therefore, since $B \rightarrow B^{\text{sh}}$ is unramified, $(g)B^{\text{sh}} = (\sqrt[e]{f})$. Since $B \rightarrow B^{\text{sh}}$ is faithfully flat, $IB^{\text{sh}} \cap B = I$ for all ideals I of B (by e.g. [4, Exercise 3.16]), so since $(g, t)B^{\text{sh}} = (\sqrt[e]{f}, t)$ is maximal, $(g, t)B^{\text{sh}} \cap B = (g, t)$ is the maximal ideal of B . Thus $\{g, t\}$ is a regular system for B .

Since t is a local equation for $\rho^{-1}C_i$, $\rho^{-1}C_i$ is regular and irreducible at y for each C_i passing through x . In particular $C_Y = \bigcup_i \rho^{-1}C_i$ is reduced, and so equals $Y_{0, \text{red}}$. Since at most two irreducible components of C meet at x , the same holds for C_Y at y , and y is a singular point on C_Y if and only if $x = f(y) \in \mathcal{S}$. This completes the proof of (b).

If $D' \in \mathcal{D}$ is the distinguished (horizontal) prime divisor running through x then there is a single irreducible component of D'_Y passing through y , whose support $D'_{Y, \text{red}}$ has local equation g at y . Thus each irreducible component of D'_Y covers D' , hence $\text{Spec } R$, hence D'_Y is horizontal. Since g is part of the regular system $\{g, t\}$ at the arbitrary closed point y we see that $D'_{Y, \text{red}}$ is regular, and since t is a local equation for an arbitrary irreducible component of C_Y passing through y , $D'_{Y, \text{red}}$ intersects each component of C_Y transversally. Thus the support of the irreducible components of D'_Y generate a set of distinguished divisors \mathcal{D}_Y for Y . This proves (c). ■

Lemma 3.2. *Suppose X is a regular noetherian scheme and L is an étale $K(X)$ -algebra that is tamely ramified along a divisor D . Then the normalization Y of X in L defines a tamely ramified cover $\rho : Y \rightarrow (X, D)$.*

Proof. Since X is regular, its connected components are integral regular schemes, hence we may assume X is integral. Since $L/K(X)$ is étale, L is a product of finite separable field extensions of $K(X)$, hence we may assume $L/K(X)$ is itself a finite separable field extension. Then the normalization Y exists, Y is normal by definition, and $\rho : Y \rightarrow X$ is finite by [26, 4.1.25]. Since Y is normal and connected it is irreducible, so Y dominates X . Let $U = X - D$, and set $V = Y \times_X U$. Since X is normal, Y is connected, and $\rho|_V$ is unramified, $\rho|_V$ is étale by [22, I.9.11] (see also [28, I.3.20]). Therefore $Y \rightarrow (X, D)$ is a tamely ramified cover. ■

The next lemma shows how distinguished divisors split in tamely ramified covers.

Lemma 3.3. *Assume the setup of (2.3). Suppose $\rho : Y \rightarrow (X, D)$ is a tamely ramified cover, where $D \in \mathcal{D}$, and $D' \in \mathcal{D}_{\mathcal{S}}$ is irreducible. Suppose $E \subset D'_{Y, \text{red}}$ is a distinguished prime divisor lying over D' as in Lemma 3.1(c), $y = E \times_Y C_Y$, and*

$x = D' \times_X C$. Then y and x are regular closed points, and the ramification (resp. inertia) degree of v_E over $v_{D'}$ equals the ramification (resp. inertia) degree of v_y over v_x .

Proof. Since we assume (2.3) and $D \in \mathcal{D}$ we have Lemma 3.1, which shows C_Y is reduced and $E \subset D'_{Y,\text{red}}$ is distinguished. Note that either $D' \cap D = \emptyset$ or $D' \subset D$. Since D' and E are distinguished and avoid the singular points of X and Y , they intersect the reduced closed fibers C and C_Y transversally, hence $x = D' \times_X C$ and $y = E \times_Y C_Y$ are regular closed points. We must show that $[\kappa(E) : \kappa(D')] = [\kappa(y) : \kappa(x)]$ and that $v_E(f) = v_y(f_0)$, where $f \in \mathcal{O}_{X,D'}$ is a local equation for D' on X and $f_0 \in \mathcal{O}_{C,x}$ is a local equation for x on C .

Since D' is horizontal and irreducible, $D' = \text{Spec } S$ for S a finite local R -algebra by [28, I.4.2], and S is a discrete valuation ring since D' is regular. The map $E \rightarrow \rho^{-1}D' \rightarrow D'$ is finite as a composition of finite morphisms, hence $E = \text{Spec } T$ for T a finite local S -algebra, again a discrete valuation ring since E is regular. Since S is a discrete valuation ring and $S \rightarrow T$ is finite, T is a free S -module of finite rank, and so $[T : S]$ is well defined. Since the generic point of E lies over that of D' , we have $\text{Frac } T = T \otimes_S \text{Frac } S$, hence $[\kappa(E) : \kappa(D')] = [T : S]$.

Let $A = \mathcal{O}_{X,x}$, $B = \mathcal{O}_{Y,y}$, let t be a local equation for C at x , and set $A_0 = A/(t)$ and $B_0 = B/(t)B$, the (reduced) local rings of the fibers through x and y , as in the proof of Lemma 3.1. Already $\kappa(x) = S \otimes_A A_0$ and $\kappa(y) = T \otimes_B B_0$ by the transversality of the intersections. Since $B_0 = B \otimes_A A_0$ we have $\kappa(y) = T \otimes_A A_0$, hence $[\kappa(y) : \kappa(x)] = [T : S] = [\kappa(E) : \kappa(D')]$ by base change.

Let $g \in A$ be defined as above. To compute the ramification degree, note that since $B \rightarrow B^{\text{sh}}$ is faithfully flat, $(g^e)B = (g^e)B^{\text{sh}} \cap B = (f)B^{\text{sh}} \cap B = (f)B$, hence $g^e = fu$ for some $u \in B^*$. Since f and g are uniformizers for $v_{D'}$ and v_E , respectively, it follows that $e(v_E/v_{D'}) = v_E(f) = e$. On the other hand, let f_0 be the image of f in A_0 , and let g_0 be the image of g in B_0 . Then f_0 cuts out the closed point x on C and g_0 cuts out y on C_Y by transversality. Thus f_0 and g_0 are uniformizers for v_x and v_y , and since $g_0^e = f_0 u_0$, where u_0 is the image of u in B_0^* , we have $e(v_y/v_x) = v_y(f_0) = e$, as desired. This completes the proof. ■

4. LIFTING COHOMOLOGY CLASSES

4.1. Let k be a field, and let C/k be a reduced connected projective curve with regular irreducible components C_1, \dots, C_m , at most two of which meet at any closed point. Denote the singular points of C by \mathcal{S} and write $\mathcal{O}_{C,\mathcal{S}}$ for the semilocal ring $\varinjlim_U \mathcal{O}_C(U)$, where U varies over (dense) open subsets of C containing \mathcal{S} . Then $\mathcal{O}_{C,\mathcal{S}}$ is a subring of the rational function ring $\kappa(C) = \prod_i \kappa(C_i)$. For each $z \in \mathcal{S} \cap C_i$, let $K_{i,z} = \text{Frac } \mathcal{O}_{C_i,z}^{\text{h}}$, a field since z is a normal point of C_i , and if $\alpha_i \in H^q(\kappa(C_i))$, let $\alpha_{i,z}$ denote the image in $H^q(K_{i,z})$.

Lemma 4.2 (Gluing). *Assume the setup of (4.1). There exists an element $\alpha \in H^q(\mathcal{O}_{C,\mathcal{S}}, \Lambda)$ that restricts to $\alpha_C = (\alpha_1, \dots, \alpha_m) \in \bigoplus_i H^q(\kappa(C_i), \Lambda)$ if and only if α_i is unramified at each $z \in \mathcal{S} \cap C_i$, and $\alpha_{i,z} = \alpha_{j,z}$ whenever $z \in C_i \cap C_j$.*

Proof. There is an exact sequence ([28, III.1.25])

$$\begin{aligned}
 0 \longrightarrow H_S^0(O_{C,S}) \longrightarrow H^0(O_{C,S}) \longrightarrow H^0(\kappa(C)) \longrightarrow H_S^1(O_{C,S}) \longrightarrow \\
 \longrightarrow H^1(O_{C,S}) \longrightarrow H^1(\kappa(C)) \longrightarrow H_S^2(O_{C,S}) \longrightarrow \\
 (*) \qquad \qquad \qquad \longrightarrow H^2(O_{C,S}) \longrightarrow H^2(\kappa(C)) \longrightarrow H_S^3(O_{C,S})
 \end{aligned}$$

where the maps into the direct sum are restrictions. Since \mathcal{S} is a disjoint union of closed points, $H_S^q(O_{C,S}) = \bigoplus_{z \in \mathcal{S}} H_z^q(O_{C,S}) = \bigoplus_{z \in \mathcal{S}} H_z^q(O_{C,z}^h)$ by excision ([28, III.1.28, p.93]). Since Λ is a smooth group scheme, $H^q(O_{C,z}^h) = H^q(\kappa(z))$, by the cohomological Hensel's lemma [28, III.3.11(a), p.116]. Since the C_k are regular and at most two of them meet at any $z \in \mathcal{S}$, we have $\text{Spec } O_{C,z}^h - \{z\} = \text{Spec } (K_{i,z} \times K_{j,z})$ for some i and j , and an “excised” exact sequence

$$\begin{aligned}
 0 \longrightarrow H_z^0(O_{C,S}) \longrightarrow H^0(\kappa(z)) \longrightarrow H^0(K_{i,z} \times K_{j,z}) \longrightarrow H_z^1(O_{C,S}) \longrightarrow \\
 \longrightarrow H^1(\kappa(z)) \longrightarrow H^1(K_{i,z} \times K_{j,z}) \longrightarrow H_z^2(O_{C,S}) \longrightarrow \\
 \longrightarrow H^2(\kappa(z)) \longrightarrow H^2(K_{i,z} \times K_{j,z}) \longrightarrow H_z^3(O_{C,S}) \longrightarrow \dots
 \end{aligned}$$

where the map $H^q(\kappa(z)) \rightarrow H^q(K_{i,z} \times K_{j,z}) = H^q(K_{i,z}) \oplus H^q(K_{j,z})$ is the diagonal map given by inflation from $\kappa(z)$ to the “local fields” $K_{i,z}$ and $K_{j,z}$. Since n is prime-to- p , the map $H^0(\kappa(z)) \rightarrow H^0(K_{i,z})$ is an isomorphism, so $H_z^0(O_{C,S}) = 0$, and for $q \geq 1$ we have short exact Witt-type sequences

$$0 \rightarrow H^q(\kappa(z)) \rightarrow H^q(K_{i,z}) \xrightarrow{\partial_z} H^{q-1}(\kappa(z), -1) \rightarrow 0$$

Thus the long exact sequence breaks up into short exact sequences

$$(4.3) \quad 0 \rightarrow H^q(\kappa(z)) \rightarrow H^q(K_{i,z} \times K_{j,z}) \rightarrow H_z^{q+1}(O_{C,S}) \rightarrow 0 \quad (q \geq 0)$$

By the compatibility of the localization sequence with the excised sequence, the map $H^q(\kappa(C_i)) \rightarrow H_z^{q+1}(O_{C,S}) \leq H_S^{q+1}(O_{C,S})$ of $(*)$ factors through $\text{res}_{\kappa(C_i)|K_{i,z}}$. Therefore an element $\alpha_C = (\alpha_1, \dots, \alpha_m) \in H^q(\kappa(C))$ maps to zero in $H_S^{q+1}(O_{C,S})$ if and only if each couple $(\alpha_{i,z}, \alpha_{j,z})$ is in the image of some $\bar{\alpha} \in H^q(\kappa(z))$; i.e., $\alpha_{i,z} = \alpha_{j,z}$, and both are unramified. Thus by the exactness of $(*)$, α_C comes from $H^q(O_{C,S})$ if and only if each α_i is unramified at each $z \in \mathcal{S} \cap C_i$, and $\alpha_{i,z} = \alpha_{j,z}$ whenever $z \in C_i \cap C_j$. ■

Suppose C is as in (4.1). Since exactly two irreducible components meet at any $z \in \mathcal{S}$ the *dual graph* G_C is defined, and consists of a vertex for each irreducible component of C and an edge for each singular point, such that an edge and a vertex are incident when the corresponding singular point lies on the corresponding irreducible component ([30, 2.23], see also [26, 10.1.48]). The (first) Betti number for G_C is $\beta_C \stackrel{\text{df}}{=} \text{rk}(H_1(G_C, \mathbb{Z})) = N + E - V$, where V, E and N are the numbers of vertices, edges, and connected components of G_C , respectively.

Lemma 4.4. *Assume the setup of (4.1). Then:*

- a) *For any integer r , $H^1(C, \mathbb{Z}/n(r)) \rightarrow H^1(O_{C,S}, \mathbb{Z}/n(r))$ is injective.*

- b) The map $H^q(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n(q-1)) \rightarrow H^q(\kappa(C), \mathbb{Z}/n(q-1))$ is injective for $q = 0, 2$, and for $q = 1$ we have

$$H^1(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n) \simeq (\mathbb{Z}/n)^{\beta_C} \oplus \Gamma$$

where $(\mathbb{Z}/n)^{\beta_C}$ is the kernel of $H^1(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n) \rightarrow H^1(\kappa(C), \mathbb{Z}/n)$, and $\Gamma \leq H^1(\kappa(C), \mathbb{Z}/n)$ is the group of tuples that glue as in Lemma 4.2.

Proof. We suppress the notation for $\Lambda = \mathbb{Z}/n(r)$. Let $z \in C - \mathcal{S}$ be a (regular) closed point, and set $U = C - \{z\}$, a dense open subset containing \mathcal{S} . The localization exact sequence is

$$\begin{aligned} 0 \longrightarrow H_z^0(C) \longrightarrow H^0(C) \longrightarrow H^0(U) \longrightarrow \cdots \\ \cdots \longrightarrow H_z^q(C) \longrightarrow H^q(C) \longrightarrow H^q(U) \longrightarrow H_z^{q+1}(C) \longrightarrow \cdots \end{aligned}$$

By excision we have an exact sequence

$$\begin{aligned} 0 \longrightarrow H_z^0(C) \longrightarrow H^0(\mathcal{O}_{C,z}^h) \longrightarrow H^0(K_z) \longrightarrow \cdots \\ \cdots \longrightarrow H_z^q(C) \longrightarrow H^q(\mathcal{O}_{C,z}^h) \longrightarrow H^q(K_z) \longrightarrow H_z^{q+1}(C) \longrightarrow \cdots \end{aligned}$$

where $K_z = \text{Frac } \mathcal{O}_{C,z}^h$. Since z is a regular point $\mathcal{O}_{C,z}^h$ is a discrete valuation ring, and by [12, Section 3.6] we may replace $H_z^{q+1}(C)$ with $H^{q-1}(\kappa(z), -1)$, and the map from $H^q(U)$, which factors through $H^q(K_z)$, is then the residue map ∂_z . We conclude $H^0(C) = H^0(U)$, and we have a long exact sequence

$$\begin{aligned} 0 \longrightarrow H^1(C) \longrightarrow H^1(U) \xrightarrow{\partial_z} H^0(\kappa(z), -1) \longrightarrow \cdots \\ (**) \quad \cdots \longrightarrow H^q(C) \longrightarrow H^q(U) \xrightarrow{\partial_z} H^{q-1}(\kappa(z), -1) \longrightarrow \cdots \end{aligned}$$

As $H^1(\mathcal{O}_{C,\mathcal{S}}) = \varinjlim_U H^1(U)$, where the limit is over dense open subsets of C containing \mathcal{S} , $H^1(C) \rightarrow H^1(\mathcal{O}_{C,\mathcal{S}})$ is injective by the exactness of the injective limit functor, proving (a).

For (b) we go back to $\Lambda = \mathbb{Z}/n$. By (*) we have an exact sequence

$$\begin{aligned} 0 \longrightarrow H^0(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n) \xrightarrow{\phi_1} H^0(\kappa(C), \mathbb{Z}/n) \xrightarrow{\phi_2} H_S^1(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n) \xrightarrow{\phi_3} \\ \xrightarrow{\phi_3} H^1(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n) \xrightarrow{\phi_4} H^1(\kappa(C), \mathbb{Z}/n) \end{aligned}$$

The groups $H^0(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n)$ and $H^0(\kappa(C), \mathbb{Z}/n)$ are finite free \mathbb{Z}/n -modules whose ranks are the number of C 's connected components N and irreducible components m , respectively. We claim $H_S^1(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n)$ is a finite free \mathbb{Z}/n -module. For by (4.3), for each $z \in \mathcal{S}$ we have an exact sequence

$$0 \longrightarrow H^0(\kappa(z), \mathbb{Z}/n) \longrightarrow H^0(K_{i,z}, \mathbb{Z}/n) \oplus H^0(K_{j,z}, \mathbb{Z}/n) \longrightarrow H_z^1(\mathcal{O}_{C,z}^h, \mathbb{Z}/n) \rightarrow 0$$

This shows $H_z^1(\mathcal{O}_{C,z}^h, \mathbb{Z}/n) \simeq \mathbb{Z}/n$, and since $H_S^1(\mathcal{O}_{C,\mathcal{S}}, \mathbb{Z}/n)$ is a finite direct sum of these groups, it is a finite free \mathbb{Z}/n -module, of rank $|\mathcal{S}|$.

The result [14, 27.1] implies that a free \mathbb{Z}/n -submodule of a \mathbb{Z}/n -module is a direct summand. Therefore we have a decomposition

$$H^0(\kappa(C), \mathbb{Z}/n) \simeq \text{im}(\phi_1) \oplus \text{im}(\phi_2)$$

and since $H^0(\kappa(C), \mathbb{Z}/n)$ is a finite free \mathbb{Z}/n -module, $\text{im}(\phi_2)$ is a finite free \mathbb{Z}/n -module by the structure theorem for finitely generated abelian groups. Similarly,

since $H_S^1(O_{C,S}, \mathbb{Z}/n)$ is a finite free \mathbb{Z}/n -module,

$$H_S^1(O_{C,S}, \mathbb{Z}/n) \simeq \text{im}(\phi_2) \oplus \text{cok}(\phi_2)$$

and since $\text{im}(\phi_2)$ and $H_S^1(O_{C,S}, \mathbb{Z}/n)$ are finite free \mathbb{Z}/n -modules, so is $\text{cok}(\phi_2)$. Since $H^1(O_{C,S}, \mathbb{Z}/n)$ is a \mathbb{Z}/n -module, $\text{cok}(\phi_2)$ is a direct summand of $H^1(O_{C,S}, \mathbb{Z}/n)$, again by [14, 27.1]. Thus we have a decomposition

$$H^1(O_{C,S}, \mathbb{Z}/n) \simeq \text{cok}(\phi_2) \oplus \text{im}(\phi_4)$$

Now we set $\Gamma = \text{im}(\phi_4)$, and compute $\text{rk}(\text{cok}(\phi_2)) = N + |\mathcal{S}| - m = \beta_C$. This proves the $q = 1$ part of (b).

The $q = 0$ case of (b) is in the proof of Lemma 4.2. Suppose $q = 2$. To show $H^2(O_{C,S}, \mu_n) \rightarrow H^2(\kappa(C), \mu_n)$ is injective, we will show $H^1(\kappa(C), \mu_n) \rightarrow H_S^2(O_{C,S}, \mu_n)$ is onto and apply the exactness of (*).

For each closed point $z \in C_1 \cap C_2 \subset \mathcal{S}$, we have a diagram

$$\begin{array}{ccccccc} H^1(\kappa(C_1), \mu_n) \oplus H^1(\kappa(C_2), \mu_n) & \longrightarrow & H_z^2(O_{C,z}^h, \mu_n) \\ \downarrow & & \parallel \\ 0 \longrightarrow H^1(\kappa(z), \mu_n) \longrightarrow H^1(\kappa(C_1)_z, \mu_n) \oplus H^1(\kappa(C_2)_z, \mu_n) & \longrightarrow & H_z^2(O_{C,z}^h, \mu_n) \longrightarrow 0 \end{array}$$

We will show that $H^1(\kappa(C_1), \mu_n) \oplus H^1(\kappa(C_2), \mu_n) \rightarrow H_z^2(O_{C,z}^h, \mu_n)$ is onto, by showing the downarrow is onto. Since z is a regular point of each C_i , each $O_{C_i,z}$ is a discrete valuation ring with residue field $\kappa(z)$ and fraction field $\kappa(C_i)$, and we have a diagram of split short exact sequences

$$\begin{array}{ccccccc} 0 \longrightarrow H^1(O_{C_i,z}, \mu_n) \longrightarrow H^1(\kappa(C_i), \mu_n) \longrightarrow H^0(\kappa(z), \mathbb{Z}/n) \longrightarrow 0 \\ \downarrow \qquad \qquad \qquad \downarrow \qquad \qquad \qquad \parallel \\ 0 \longrightarrow H^1(\widehat{O}_{C_i,z}, \mu_n) \longrightarrow H^1(\kappa(C_i)_z, \mu_n) \longrightarrow H^0(\kappa(z), \mathbb{Z}/n) \longrightarrow 0 \end{array}$$

To show the middle downarrow is onto it suffices (by a standard diagram chase) to prove that the left downarrow is onto. Since $\widehat{O}_{C_i,z}$ is henselian $H^1(\widehat{O}_{C_i,z}, \mu_n) = H^1(\kappa(z), \mu_n)$, and by Kummer theory and Hilbert 90 we have $H^1(O_{C_i,z}, \mu_n) = O_{C_i,z}^*/n$ and $H^1(\kappa(z), \mu_n) = \kappa(z)^*/n$. Since $O_{C_i,z} \rightarrow \kappa(z)$ is onto and $O_{C_i,z}$ is local, the induced map $O_{C_i,z}^* \rightarrow \kappa(z)^*$ is onto, hence $H^1(O_{C_i,z}, \mu_n)$ maps onto $H^1(\kappa(z), \mu_n)$. We conclude $H^1(\kappa(C_i), \mu_n) \rightarrow H^1(\kappa(C_i)_z, \mu_n)$ is onto. Now each map $H^1(\kappa(C_1), \mu_n) \oplus H^1(\kappa(C_2), \mu_n) \rightarrow H_z^2(O_{C,z}^h, \mu_n)$ is onto.

Suppose $(b_z) \in H_S^2(O_{C,S}, \mu_n) = \bigoplus_{\mathcal{S}} H_z^2(O_{C,z}^h, \mu_n)$. We have just seen that for each closed point $z \in C_i \cap C_j$ there exists a pair $([a_{i,z} t_{i,z}^{e_i}], [a_{j,z} t_{j,z}^{e_j}]) \in \kappa(C_i)^*/n \oplus \kappa(C_j)^*/n$ mapping to b_z , for z -units $a_{k,z} \in O_{C_k,z}^*$, z -uniformizers $t_{k,z} \in \kappa(C_k)$, and integers e_k , for $k = i, j$. Let $v_{k,z}$ be the discrete valuation on $\kappa(C_k)$ determined by z . By standard approximation (e.g. [25, XII.1.2]) there exist elements $a_k, t_k \in \kappa(C_k)$ such that

$$v_{k,z}(a_k - a_{k,z}) > 0 \quad \text{and} \quad v_{k,z}(t_k - t_{k,z}) > 1 \quad \text{for all } z.$$

The image of $a_k t_k^{e_k}$ in $\kappa(C_k)_z^*/n$ is $[a_{k,z} t_{k,z}^{e_k}]$. Therefore the m -tuple

$$([a_k t_k^{e_k}]) \in H^1(\kappa(C), \mu_n)$$

maps to (b_z) . This proves the induced map $H^1(\kappa(C), \mu_n) \rightarrow H_S^2(\mathcal{O}_{C,S}, \mu_n)$ is onto, and completes the proof. \blacksquare

We will soon need the following technical lemma in order to replace X_0 with C .

Lemma 4.5. *Suppose A is a noetherian ring. Then $(\text{Frac } A)_{\text{red}} = \text{Frac } (A_{\text{red}})$ if and only if A has no embedded primes.*

Proof. By definition $\text{Frac } A = S^{-1}A$, where $S = A - \bigcup_{\text{Ass } A} \mathfrak{p}$, and $S^{-1}N_A = N_{S^{-1}A}$ by [4, 3.12], hence $S^{-1}(A_{\text{red}}) = (S^{-1}A)_{\text{red}}$. It remains to show that $S^{-1}(A_{\text{red}}) = \text{Frac } (A_{\text{red}})$ if and only if A has no embedded primes. By $S^{-1}(A_{\text{red}})$ of course we mean $f(S)^{-1}(A_{\text{red}})$, where $f : A \rightarrow A_{\text{red}}$. This localization equals the localization with respect to the multiplicative set T , where T is the saturation of $f(S)$ in A_{red} , and this is the complement of the union of prime ideals of A_{red} that don't meet $f(S)$ by [4, Exercise 3.7]. Thus $S^{-1}(A_{\text{red}}) = \text{Frac } (A_{\text{red}})$ if and only if the union of the primes of A_{red} that don't meet $f(S)$ equals the union of the associated primes of A_{red} , which are just the minimal primes of A_{red} . But A and A_{red} have identical underlying topological spaces, and the primes of A_{red} that don't meet $f(S)$ correspond to the primes of A that don't meet S , i.e., the associated primes. These correspond to the minimal primes of A_{red} if and only if the associated primes of A are the minimal primes of A , i.e., A has no embedded primes. \square

Theorem 4.6. *Assume the setup of (2.3), with X connected. Then for $q \geq 0$ there is a map*

$$\lambda : H^q(\mathcal{O}_{C,S}, \Lambda) \rightarrow H^q(K(X), \Lambda)$$

and a commutative diagram

$$(4.7) \quad \begin{array}{ccc} H^q(\mathcal{O}_{C,S}, \Lambda) & \xrightarrow{\lambda} & H^q(K(X), \Lambda) \\ \oplus \text{res}_i \downarrow & & \downarrow \oplus_i \text{res}_i \\ H^q(\kappa(C), \Lambda) & \xrightarrow{\text{inf}} & \bigoplus_i H^q(K(X)_{C_i}, \Lambda) \end{array}$$

such that if $\alpha_0 \in H^q(\mathcal{O}_{C,S}, \Lambda)$ and $\alpha = \lambda(\alpha_0)$ then:

- a) α is defined at the generic points of C_i , and $\alpha(C_i) = \text{res}_i(\alpha_0)$.
- b) The ramification locus of α (on X) is contained in \mathcal{D}_S .
- c) If $D \in \mathcal{D}_S$ is prime and $z = D \cap C$, then $\partial_D \cdot \lambda = \inf_{\kappa(z)|\kappa(D)} \partial_z$.
- d) If α_0 is unramified at a closed point z , and D is any (horizontal) prime lying over z , then α is unramified at D , and has value $\alpha(D) = \inf_{\kappa(z)|\kappa(D)} (\alpha_0(z))$.

Proof. Let D_0 be an effective divisor on C that avoids \mathcal{S} , let $D \in \mathcal{D}_S$ be the distinguished lift of D_0 , set $U = X - D$, and set $U_0 = C - D_0$. Since X and D are regular and D has pure codimension 1, we have $H^0(X) \simeq H^0(U)$, and an exact Gysin sequence

$$0 \rightarrow H^1(X) \rightarrow H^1(U) \xrightarrow{\partial_D} H^0(D, -1) \rightarrow H^2(X) \rightarrow \dots$$

by Gabber's absolute purity theorem ([15, Theorem 2.1.1]) and the standard construction of the Gysin sequence ([12, Section 3.2]). (Note that the result in [15] is stated for the $\Lambda = \mathbb{Z}/n$ case only, but the result holds in general since the sheaves $\mathcal{H}_D^q(X)$ and $\mathcal{H}_D^q(X, \mathbb{Z}/n)$ are locally isomorphic, and the morphism $i^*\Lambda(-1) \rightarrow$

$\mathcal{H}_D^2(X)$ is canonical.) We use the notation ∂_D since this map is compatible with the one defined above on $H^q(K(X))$ when D is prime.

We may replace X_0 by $C = X_{0,\text{red}}$ in the cohomological computations below since Λ is finite and n is prime-to- p , by [28, V.2.4(c)] (see also [28, II.3.11]). To substitute $\mathcal{O}_{C,S}$ and $\kappa(C)$ for $\mathcal{O}_{X_0,S}$ and $\kappa(X_0)$ we must check that the former are the canonical reduced quotients of the latter. But the ring $\mathcal{O}_{X_0,S}$ can be obtained by localizing some affine open subset $\text{Spec } A_0$ containing \mathcal{S} (which exists since X_0/k is projective) with respect to the multiplicative set $T = A_0 - \bigcup_{\mathcal{S}} \mathfrak{m}_z$. Since $\mathcal{O}_{C,S}$ is obtained by localizing $A_{0,\text{red}}$ with respect to the image of T in $A_{0,\text{red}}$, we have $\mathcal{O}_{C,S} = (\mathcal{O}_{X_0,S})_{\text{red}}$ since the formation of the nilradical commutes with localization (see e.g. [4, 3.12]).

To show $\kappa(C) = \kappa(X_0)_{\text{red}}$ it suffices to show X_0 has no embedded points by Lemma 4.5. But if z is any closed point of X then $\mathcal{O}_{X,z}$ is a regular local ring, and a local equation for the closed fiber $\mathcal{O}_{X,z} \otimes_R k$ passing through z is given by the uniformizer p in R . Since $\mathcal{O}_{X,z}$ is factorial and at most two components of X_0 pass through z we have $p = \pi_1^{e_1} \pi_2^{e_2}$ for primes π_i and numbers $e_i \geq 0$. The associated primes of $\mathcal{O}_{X,z}/(\pi_1^{e_1} \pi_2^{e_2})$ are evidently just the (π_i) , which shows X_0 has no embedded point at z .

Since D_0 is a disjoint union of regular closed points, by $(**)$ and the work that immediately precedes it we have $H^0(C) \simeq H^0(U_0)$ and an exact sequence

$$0 \longrightarrow H^1(C) \longrightarrow H^1(U_0) \xrightarrow{\partial_{D_0}} H^0(\kappa(D_0), -1) \longrightarrow H^2(C) \longrightarrow \dots$$

Thus we have a commutative ladder

$$\begin{array}{ccccccc} 0 & \longrightarrow & H^1(X) & \longrightarrow & H^1(U) & \xrightarrow{\partial_D} & H^0(D, -1) \longrightarrow H^2(X) \longrightarrow \dots \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & H^1(C) & \longrightarrow & H^1(U_0) & \xrightarrow{\partial_{D_0}} & H^0(D_0, -1) \longrightarrow H^2(C) \longrightarrow \dots \end{array}$$

Since R is complete, $H^q(X) \rightarrow H^q(C)$ and $H^q(D, -1) \rightarrow H^q(D_0, -1)$ are isomorphisms for $q \geq 0$ by proper base change ([28, VI.Corollary 2.7]). Therefore, in light of the isomorphisms in degree zero and the 5-lemma in degree $q \geq 1$, we obtain isomorphisms

$$H^q(U) \xrightarrow{\sim} H^q(U_0)$$

for $q \geq 0$. Let \tilde{U} denote the inverse limit over all these open sets U (this is a scheme by [20, 8.2.3]). Then $H^q(\tilde{U})$ is the direct limit of the $H^q(U)$ by [28, III.1.16], and since the direct limit functor is exact we have an isomorphism $H^q(\tilde{U}) \xrightarrow{\sim} H^q(\mathcal{O}_{C,S})$. Composing the inverse with $H^q(\tilde{U}) \rightarrow H^q(K(X))$ yields our lift

$$\lambda : H^q(\mathcal{O}_{C,S}) \longrightarrow H^q(K(X))$$

The commutative diagram (4.7) follows by applying cohomology to the diagram

$$\begin{array}{ccccc} \text{Spec } \mathcal{O}_{C,S} & \longrightarrow & \tilde{U} & \longleftarrow & \text{Spec } K(X) \\ \uparrow & & \uparrow & & \uparrow \\ \text{Spec } \kappa(C_i) & \longrightarrow & \text{Spec } \mathcal{O}_{K(X)_{C_i}} & \longleftarrow & \text{Spec } K(X)_{C_i} \end{array}$$

incorporating the isomorphisms induced by the upper and lower left horizontal arrows. If $\alpha_0 \in H^q(\mathcal{O}_{C,\mathcal{S}})$ and $\alpha = \lambda(\alpha_0)$, then since \tilde{U} contains \mathcal{S} and the generic points of the C_i , α is defined at these points, and the formula $\alpha(C_i) = \text{res}_{\mathcal{O}_{C,\mathcal{S}}|\kappa(C_i)}(\alpha_0)$ follows immediately from (4.7), proving (a).

If D is a horizontal prime divisor not in $\mathcal{D}_{\mathcal{S}}$, then the generic point $\text{Spec } \kappa(D)$ is contained in \tilde{U} , hence the map $H^q(\tilde{U}) \rightarrow H^q(K(X)_D)$ factors through $H^q(\mathcal{O}_{K(X)_D})$, which shows $\partial_D \cdot \lambda = 0$. Thus the ramification locus of any element in the image of λ must be contained in $\mathcal{D}_{\mathcal{S}}$, proving (b). Now if $D \in \mathcal{D}_{\mathcal{S}}$ is prime and $z = D \cap C$ then D is the prime spectrum of a complete local ring with residue field $\kappa(z)$, and the isomorphism

$$H^{q-1}(D, -1) \xrightarrow{\sim} H^{q-1}(z, -1) = H^{q-1}(\kappa(z), -1)$$

is the standard identification. Thus the formula $\partial_D \cdot \lambda = \inf_{\kappa(z)|\kappa(D)} \cdot \partial_z$ is immediate by the commutative ladder of Gysin sequences above, proving (c).

Suppose $\alpha = \lambda(\alpha_0)$ has ramification locus D_α , then $D_\alpha \in \mathcal{D}_{\mathcal{S}}$. Set $U = X - D_\alpha$. If α_0 is unramified at a point z , then α is unramified at every prime divisor D lying over z . For if $D \in \mathcal{D}_{\mathcal{S}}$ then $\partial_D(\alpha) = \inf(\partial_z(\alpha_0))$ by the formula just proved, and if $D \notin \mathcal{D}_{\mathcal{S}}$ then $\partial_D(\alpha) = 0$ since $D_\alpha \in \mathcal{D}_{\mathcal{S}}$. Thus if α_0 is unramified at z , and D is a prime divisor lying over z , then U contains D . The maps $z = \text{Spec } \kappa(z) \rightarrow U_0$ and $D \rightarrow U$ then induce a commutative diagram

$$\begin{array}{ccccc} H^q(U) & \xrightarrow{\text{res}} & H^q(D) & \xrightarrow{\text{res}} & H^q(\kappa(D)) \\ \downarrow \text{res} & & \downarrow \text{res} & \nearrow \text{inf} & \\ H^q(U_0) & \xrightarrow{\text{res}} & H^q(z) & & \end{array}$$

Both vertical down-arrows are isomorphisms by proper base change. The inverse of the left one is λ by definition, and the composition of the inverse of the right one and the restriction $H^q(D) \rightarrow H^q(\kappa(D))$ is inflation, as shown. Since $\kappa(D)$ is complete, the top composition of horizontal restrictions factors through the restriction $H^q(U) \rightarrow H^q(\hat{\mathcal{O}}_{X,D})$ and the bottom factors through the restriction $H^q(U_0) \rightarrow H^q(\hat{\mathcal{O}}_{C,z})$. Since these are restriction maps, the images of α and α_0 are the values $\alpha(D)$ and $\alpha_0(z)$. We conclude $\inf_{\kappa(z)|\kappa(D)}(\alpha_0(z)) = \alpha(D)$, as in (d). ■

4.8. By weak approximation [34, Lemma] there exists a $\pi \in K(X)$ such that

$$\text{div } \pi = C + E$$

where E contains no components of C , and avoids any finite set of points \mathcal{N} . We fix such a π for \mathcal{N} containing \mathcal{S} . For each i , the choice of π determines a noncanonical “Witt” isomorphism

$$H^q(\kappa(C_i)) \oplus H^{q-1}(\kappa(C_i), -1) \xrightarrow{\sim} H^q(K(X)_{C_i})$$

Taking (α, θ) to $\alpha + (\pi) \cdot \theta$, where α and θ are inflated from $\kappa(C_i)$ to $K(X)_{C_i}$, (π) is the image of π in $H^1(K(X)_{C_i}, \mu_n)$, and $(\pi) \cdot \theta$ is the cup product. Although we cannot in general lift all of $\bigoplus_i H^q(K(X)_{C_i})$ to $H^q(K(X))$, we can now prove the following.

Corollary 4.9. *Let (π) denote the image of π in $H^1(K(X), \mu_n)$. The choice of \mathcal{D}_S and π determines a homomorphism for $q \geq 1$,*

$$\begin{aligned} \lambda : H^q(\mathcal{O}_{C,S}, \Lambda) \oplus H^{q-1}(\mathcal{O}_{C,S}, \Lambda(-1)) &\longrightarrow H^q(K(X), \Lambda) \\ (\alpha_0, \theta_0) &\longmapsto \lambda(\alpha_0) + (\pi) \cdot \lambda(\theta_0) \end{aligned}$$

such that $(\bigoplus_i \text{res}_{K(X)|K(X)_{C_i}}) \cdot \lambda = \bigoplus_i (\inf_{\kappa(C_i)|K(X)_{C_i}} \cdot \text{res}_{\mathcal{O}_{C,S}|\kappa(C_i)})$.

Proof. This is an immediate consequence of Theorem 4.6. ■

Remark 4.10. a) Theorem 4.6 and Corollary 4.9 apply with obvious amendments to the case where X is not connected. For if $X = \coprod_k X_k$ is a decomposition into connected components, then $K(X) = \prod_k K(X_k)$, $X_0 = \coprod_k (X_k)_0$, $\mathcal{O}_{X_0,S} = \prod_k \mathcal{O}_{(X_k)_0, S_k}$ (where $S_k = S \cap X_k$), all of the cohomology groups break up into direct sums, and we define the map λ to be the direct sum of the maps on the summands. This will come up in the next section.

b) If X is smooth, then S is empty, and $\mathcal{O}_{C,S} = \kappa(C)$. By Witt's theorem we have $H^q(K(X)_C, \Lambda) \simeq H^q(\kappa(C), \Lambda) \oplus H^{q-1}(\kappa(C), \Lambda(-1))$, and we obtain a map

$$\lambda : H^q(K(X)_C, \Lambda) \longrightarrow H^q(K(X), \Lambda)$$

that splits the restriction map. This is the map of [10].

4.11. Completely split characters. In [30, 2.1] Saito defines a completely split covering of a noetherian scheme X to be a finite étale cover $Y \rightarrow X$ such that $Y \times_X \text{Spec } \kappa(x) = \coprod \text{Spec } \kappa(x)$, for all closed points $x \in X$. We abuse Saito's terminology (see Remark(4.13) below) and in the setup of (2.3) denote by $H_{\text{cs}}^1(C, \mathbb{Z}/n)$ the kernel of the map $H^1(\mathcal{O}_{C,S}, \mathbb{Z}/n) \rightarrow H^1(\kappa(C), \mathbb{Z}/n)$ in Lemma 4.4. If $\beta \in H_{\text{cs}}^1(C, \mathbb{Z}/n)$ then $\partial_z(\beta) = 0$ for all closed points $z \in C - S$ since ∂_z factors through $\kappa(C)_z$. Therefore β is defined on C , hence $H_{\text{cs}}^1(C, \mathbb{Z}/n) \leq H^1(C, \mathbb{Z}/n)$. Let $H_{\text{cs}}^1(X, \mathbb{Z}/n)$ denote the preimage of $H_{\text{cs}}^1(C, \mathbb{Z}/n)$ under the proper base change isomorphism.

Proposition 4.12. *Assume the setup of (2.3). Then elements of $H_{\text{cs}}^1(C, \mathbb{Z}/n)$ are trivial at all points of C , and the nontrivial elements of $H_{\text{cs}}^1(X, \mathbb{Z}/n)$ are trivial at all points of X except for the generic point $\text{Spec } K(X)$, where they are nontrivial.*

Proof. Suppose $\beta_0 \in H_{\text{cs}}^1(C)$. Then β_0 is trivial at each generic point of C by definition of $H_{\text{cs}}^1(C)$. If $z \in C$ is a closed point lying on the irreducible component C_i then the map $H^1(C) \rightarrow H^1(\kappa(z))$ factors through $H^1(C_i)$. Since C_i is regular the map $H^1(C_i) \rightarrow H^1(\kappa(C_i))$ is injective by purity, and consequently $\beta_0(z) = 0$ by definition. Thus the elements of $H_{\text{cs}}^1(C)$ are trivial at all points of C .

Suppose $\beta = \lambda(\beta_0) \in H_{\text{cs}}^1(X)$. If $x \in X$ is a generic point of some irreducible component C_i of C then the image of β in $H^1(\kappa(C_i))$ is zero since the map $H_{\text{cs}}^1(X, \mathbb{Z}/n) \rightarrow H^1(\kappa(C_i))$ factors through $H_{\text{cs}}^1(C)$. If x is the generic point of a horizontal divisor D with closed point z then $\beta(D) = \inf_{\kappa(z)|\kappa(D)}(\beta_0(z))$ by Theorem 4.6(d), and this is zero since $\beta_0(z) = 0$. If z is a closed point of X then z is on C , and the map $H_{\text{cs}}^1(X) \rightarrow H^1(\kappa(z))$ factors through $H_{\text{cs}}^1(C)$, hence β is trivial at z . Finally, since X is regular the map $H^1(X) \rightarrow H^1(K(X))$ is injective by purity, hence β is nontrivial at the generic point of X . □

Remark 4.13. Proposition 4.12 shows the elements of $H_{\text{cs}}^1(C, \mathbb{Z}/n)$ are completely split in the sense of [30]. However, in the general case our $H_{\text{cs}}^1(C, \mathbb{Z}/n)$ does not account for elements that are split at every closed point but nontrivial at generic points of C . This is not an issue if k is finite as shown by Saito in [30, Theorem 2.4], since then the C_i have no nontrivial completely split covers, essentially by Cebotarev's density theorem (see [29, Lemma 1.7]).

5. INDEX CALCULATION IN THE BRAUER GROUP

5.1. Cyclic Covers. If U is any scheme, and \bar{u} is a geometric point, the fiber functor defines a category equivalence between (finite) étale covers of U and finite continuous $\pi_1(U, \bar{u})$ -sets, yielding a canonical isomorphism

$$H^1(U, \mathbb{Z}/n) \simeq H^1(\pi_1(U, \bar{u}), \mathbb{Z}/n) = \text{Hom}_{\text{cont}}(\pi_1(U, \bar{u}), \mathbb{Z}/n)$$

(see [13, I.2.11]). If $\theta \in H^1(U, \mathbb{Z}/n)$, we will write $U[\theta]$ for the finite cyclic étale cover determined by θ . If $U = \text{Spec } A$ is affine, we will write $A[\theta]$ for the corresponding ring, or $A(\theta)$ if A is a field. If U is a connected normal scheme, and $\theta \in H^1(U, \mathbb{Z}/n)$ has order m , then $U[\theta]$ is a disjoint sum of n/m connected \mathbb{Z}/m -Galois covers of U .

Lemma 5.2. Assume the setup of (2.3). Let $\theta_0 \in H^1(\mathcal{O}_{C,S}, \mathbb{Z}/n)$ be a (tamely ramified) character with ramification divisor D_0 on C . Then the (tame) ramification divisor of $\theta = \lambda(\theta_0)$ is the distinguished lift $D \in \mathcal{D}_S$ of D_0 on X , and θ defines a tamely ramified cover $\rho : Y \rightarrow (X, D)$ as in Lemma 3.2. Restriction to C yields a tamely ramified cover $\rho_0 : C_Y \rightarrow (C, D_0)$ such that $\mathcal{O}_{C_Y, S_Y} = \mathcal{O}_{C,S}[\theta_0]$, and the reduced closed fiber C_Y of Y is the normalization of C in $\kappa(C)(\theta_0)$.

Proof. The lift θ is tamely ramified with respect to D by Theorem 4.6. Let Y be the normalization of X in $L = K(X)(\theta)$. Since D is in \mathcal{D}_S and X/R satisfies the setup of (2.3), by Lemma 3.2 $\rho : Y \rightarrow (X, D)$ is a tamely ramified cover, Y/R is a regular relative curve with reduced closed fiber C_Y , the irreducible components of C_Y are regular with singular points S_Y , and D_Y is in \mathcal{D}_{Y, S_Y} .

Let $U = X - D$, $V = U \times_X Y$, $U_0 = U \times_X X_0$ and $V_0 = V \times_Y Y_0$. Then $\theta_0 \in H^1(U_0)$. We have $H^1(U) \simeq \text{Hom}(\pi_1(U), \mathbb{Z}/n)$ and $H^1(U_0) \simeq \text{Hom}(\pi_1(U_0), \mathbb{Z}/n)$, and the restriction map $\text{res} : H^1(U) \rightarrow H^1(U_0)$, which sends θ to θ_0 , is induced by the natural map $\pi_1(U_0) \rightarrow \pi_1(U)$, which is induced on covers by $W \mapsto W \times_U U_0$. Therefore $V_0 = U_0[\theta_0]$.

We show $\mathcal{O}_{C_Y, S_Y} = \mathcal{O}_{C,S}[\theta_0]$. If $U_0 = \text{Spec } A_0 \subset C - D_0$ is a dense affine open subset of C containing S , then its preimage in C_Y is a dense affine open subset $V_0 = \text{Spec } B_0$ containing S_Y . By base change we have $B_0 = A_0[\theta_0]$, and $S^{-1}B_0 = \mathcal{O}_{C,S}[\theta_0]$, where $S = A_0 - (\bigcup_{x \in S} \mathfrak{m}_x)$ is the multiplicative set defining $\mathcal{O}_{C,S}$. Since $S_Y = \rho^{-1}S$, the saturation T of S in B_0 is $T = B_0 - (\bigcup_{y \in S_Y} \mathfrak{m}_y)$, which shows $S^{-1}B_0 = \mathcal{O}_{C_Y, S_Y}$. Therefore $\mathcal{O}_{C_Y, S_Y} = \mathcal{O}_{C,S}[\theta_0]$, and it follows immediately that $\mathcal{O}_{C_Y, S_Y} = \mathcal{O}_{C,S}[\theta_0]$.

The map $C_Y \rightarrow C$ is finite by base change, and since each irreducible component of C_Y is regular by Lemma 3.1 each irreducible component of C_Y is the normalization of a component of C in a field extension which is a direct factor of $\kappa(C_Y)$. Equivalently, C_Y is the normalization of C in $\kappa(C)(\theta_0)$, by [18, 6.3.7]. This completes the proof.

■

5.3. Index. Assume the setup of (2.3) with $R = \mathbb{Z}_p$ and $\Lambda = \mu_n$. Fix $\alpha_C \in H^2(\mathcal{O}_{C,S})$ and $\theta_C \in \Gamma \leq H^1(\mathcal{O}_{C,S}, -1)$, and write

$$\alpha_C = (\alpha_{C_1}, \dots, \alpha_{C_m}) \in H^2(\kappa(C))$$

$$\theta_C = (\theta_{C_1}, \dots, \theta_{C_m}) \in H^1(\kappa(C), -1)$$

as per Lemma 4.4(b). Suppressing the inflation maps, we form the element

$$\gamma_C = \alpha_C + (\pi) \cdot \theta_C = (\alpha_{C_1} + (\pi) \cdot \theta_{C_1}, \dots, \alpha_{C_m} + (\pi) \cdot \theta_{C_m}) \in \bigoplus_{i=1}^m H^2(K(X)_{C_i})$$

where π is as in (4.8). Let $\theta = \lambda(\theta_C)$, and let Y be the normalization of X in $K(X)(\theta)$, as in Lemma 5.2. Then C_Y is the reduced closed fiber of Y , and we write $C_{i,Y}$ for the preimage of C_i , so that $\kappa(C_{i,Y}) = \kappa(C_i)(\theta_{C_i})$. Thus

$$\alpha_{C_Y} = (\alpha_{C_{1,Y}}, \dots, \alpha_{C_{m,Y}}) \in \bigoplus_{i=1}^m H^2(\kappa(C_{i,Y}))$$

Note $\kappa(C_{i,Y})$ is a product of the function fields of the irreducible components of $C_{i,Y}$. By the (well-known) Nakayama-Witt index formula,

$$\text{ind}(\alpha_{C_i} + (\pi) \cdot \theta_{C_i}) = |\theta_{C_i}| \cdot \text{ind}(\alpha_{C_{i,Y}})$$

We have $|\theta_C| = \text{lcm}_i\{|\theta_{C_i}|\}$ by Lemma 4.4(b). We now *define*

$$\text{ind}(\alpha_{C_Y}) \stackrel{\text{df}}{=} \text{lcm}_i\{\text{ind}(\alpha_{C_{i,Y}})\}$$

$$\text{ind}(\gamma_C) \stackrel{\text{df}}{=} |\theta_C| \cdot \text{ind}(\alpha_{C_Y})$$

Theorem 5.4. Assume the setup of (2.3) with $R = \mathbb{Z}_p$. Let $\Gamma \leq H^1(\mathcal{O}_{C,S})$ be as in Lemma 4.4(b). Then the map $\lambda : H^2(\mathcal{O}_{C,S}) \oplus \Gamma \rightarrow H^2(K(X))$ preserves index.

Proof. We may assume X is connected. We identify Γ with the image of $H^1(\mathcal{O}_{C,S}, -1)$ in $H^1(\kappa(C), -1)$, as in Lemma 4.4(b), and adopt the notation of (5.3). Set $\gamma = \lambda(\gamma_C)$, $\alpha = \gamma(\alpha_C)$, and $\theta = \lambda(\theta_C)$, so that $\gamma = \alpha + (\pi) \cdot \theta$ as in Corollary 4.9. By restricting to connected components if necessary we may assume that Y is connected, hence that $K(Y)$ is a field. Even so, the cyclic-Galois étale $\kappa(C_i)$ -algebra $\kappa(C_{i,Y})$ may not be a field. Since θ_C is in $H^1(\mathcal{O}_{C,S}, -1)$ the ramification divisor D_0 of θ_C avoids \mathcal{S} , and the distinguished lift $D \in \mathcal{D}_{\mathcal{S}}$ of D_0 is the ramification divisor of θ by Theorem 4.6. By Lemma 5.2 $\rho : Y \rightarrow (X, D)$ is a cyclic tamely ramified cover, and by Lemma 3.1 Y/R satisfies the properties of (2.3), with reduced closed fiber C_Y , $\mathcal{S}_Y = \rho^{-1}\mathcal{S}$ the singular points of C_Y , and \mathcal{D}_Y generated by $D_{Y,\text{red}}$ and the preimages of the other distinguished divisors of X .

The index of γ cannot exceed $|\theta|\text{ind}(\alpha_Y)$. For if $M/K(Y)$ is a separable maximal subfield of the division algebra associated with α_Y , then $M/K(X)$ splits γ , and has degree $|\theta|\text{ind}(\alpha_Y)$. Since in our case $|\theta| = [K(Y) : K(X)] = [\kappa(C_Y) : \kappa(C)] = |\theta_C|$, to prove the theorem it is enough to prove $\text{ind}(\alpha_Y) = \text{ind}(\alpha_{C_Y})$.

By Lemma 5.2 $\mathcal{O}_{C,S}[\theta_C] = \mathcal{O}_{C_Y, \mathcal{S}_Y}$. Each $\kappa(C_{i,Y}) = \kappa(C_i)(\theta_{C_i})$ is a product of global fields, and by class field theory the division algebra associated with the restriction of $\alpha_{C_{i,Y}}$ to each field component is cyclic. Since α_{C_Y} is in $H^2(\mathcal{O}_{C_Y, \mathcal{S}_Y})$ (by Lemma 5.2 and Lemma 4.2), $\alpha_{C_{i,Y}}$ is unramified at \mathcal{S}_Y . By Grunwald-Wang's

theorem there exists a tuple $\psi_{C_Y} = (\psi_{C_{1,Y}}, \dots, \psi_{C_{m,Y}}) \in \bigoplus_i H^1(\kappa(C_{i,Y}), -1)$ such that $|\psi_{C_{i,Y}}| = \text{ind}(\alpha_{C_{i,Y}})$, $\kappa(\psi_{C_{i,Y}})$ splits $\alpha_{C_{i,Y}}$, and such that the $\psi_{C_{i,Y}}$ are unramified and equal at the local fields defined by the singular points \mathcal{S}_Y . Then ψ_{C_Y} comes from $H^1(\mathcal{O}_{C_Y, \mathcal{S}_Y}, -1)$ by Lemma 4.2, and $|\psi_{C_Y}| = \text{ind}(\alpha_{C_Y})$.

By Theorem 4.6 (and Remark 4.10(a) if Y is not connected) we have a map $\lambda_Y : H^1(\mathcal{O}_{C_Y, \mathcal{S}_Y}, -1) \rightarrow H^1(K(Y), -1)$. Since the distinguished divisors on Y are the (reduced) preimages of those on X , λ_Y is compatible with λ and the residue maps. Set $\psi = \lambda_Y(\psi_{C_Y})$, and let D_ψ denote the distinguished lift of the ramification divisor $D_{\psi_{C_Y}}$ of ψ_{C_Y} on C_Y . By Lemma 5.2, ψ determines a cyclic tamely ramified cover $\sigma : Z \rightarrow (Y, D_\psi)$ with reduced closed fiber C_Z , such that C_Z is cyclic and tamely ramified over $(C_Y, D_{\psi_{C_Y}})$, inducing ψ_{C_Y} . Since $\kappa(C_{i,Z})$ splits $\alpha_{C_{i,Y}}$, $\alpha_{C_Z} = 0$.

Again we may assume Z is connected. By construction, $[K(Z) : K(Y)] = |\psi| = |\psi_{C_Y}| = [\kappa(C_Z) : \kappa(C_Y)] = \text{ind}(\alpha_{C_Y})$. By (4.7) we have $\text{ind}(\alpha_{C_Y}) \leq \text{ind}(\alpha_Y)$, and it remains to show $\alpha_Z = 0$. It is then enough to show α_Z is unramified with respect to all Weil divisors on Z , by [10, Lemma 3.5].

Let $D' = D \cup \rho(D_\psi)$. Since $D_\psi \in \mathcal{D}_{Y, \mathcal{S}_Y}$, $\rho(D_\psi) \in \mathcal{D}_S$, and the composition $\rho' : Z \rightarrow (X, D')$ is a tamely ramified cover. Since X is regular, ρ' is (finite and) flat by Lemma 3.1, and so the image of any prime divisor J of Z is a prime divisor $\rho'(J) = I$ of X . By the functoriality of the residue maps, α_Z can only be ramified at prime divisors lying over irreducible components of D_α . By Theorem 4.6(a) D_α is in \mathcal{D}_S , and the divisors of Z lying over D_α are distinguished by Lemma 3.1(c). Thus it is enough to show that α_Z is unramified at these distinguished divisors. Clearly we may assume D_α is irreducible.

Let $E \subset Z$ be a (distinguished) prime divisor lying over $D_\alpha \in \mathcal{D}_S$. Since $D_\alpha \cap D' = \emptyset$ or $D_\alpha \subset D'$, we have $e(v_E/v_{D_\alpha}) = e(v_{E_0}/v_{D_{\alpha_C}}) = e$ for some $e \geq 1$, by Lemma 3.3. By Lemma 4.6 and the functorial behavior of the residue and restriction maps, we have a commutative diagram

$$\begin{array}{ccccc}
H^2(\mathcal{O}_{C,S}) & \xrightarrow{\lambda} & H^2(K(X)) & \xrightarrow{\text{res}} & H^2(K(Z)) \\
\partial_{D_{\alpha_C}} \downarrow & & \partial_{D_\alpha} \downarrow & & \downarrow \partial_E \\
H^1(\kappa(D_{\alpha_C}), -1) & \xrightarrow{1} & H^1(\kappa(D_\alpha), -1) & \xrightarrow{e \cdot \text{res}} & H^1(\kappa(E), -1) \\
& & \uparrow \text{inf} & & \uparrow \text{inf} \\
& & H^1(\kappa(D_{\alpha_C}), -1) & \xrightarrow{e \cdot \text{res}} & H^1(\kappa(E_0), -1) \\
\partial_{D_{\alpha_C}} \uparrow & & \uparrow & & \uparrow \partial_{E_0} \\
H^2(\mathcal{O}_{C,S}) & \xrightarrow{\text{res}} & H^2(\mathcal{O}_{C_Z, \mathcal{S}_Z}) & &
\end{array}$$

1 2 3 4

Since $\alpha = \lambda(\alpha_C)$, $\partial_E(\alpha_Z) = e \cdot (\partial_{D_{\alpha_C}}(\alpha_C))_{\kappa(E)}$ by squares (1) and (2), and by square (4), $\partial_{E_0}(\alpha_{C_Z}) = e \cdot (\partial_{D_{\alpha_C}}(\alpha_C))_{\kappa(E_0)}$. Therefore $\partial_E(\alpha_Z) = \text{inf}_{\kappa(E_0)|\kappa(E)}(\partial_{E_0}(\alpha_{C_Z}))$ by square (3). Since $\alpha_{C_Z} = 0$, we conclude $\partial_E(\alpha_Z) = 0$, as desired. This proves the theorem. ■

6. NONCROSSED PRODUCTS AND INDECOMPOSABLE DIVISION ALGEBRAS

A (finite-dimensional) division algebra D central over a field F is called a *noncrossed product* if it has no Galois maximal subfield. Its algebra structure then cannot be given by a Galois 2-cocycle, counter to almost all known division algebra constructions (see [24] for a construction of a noncrossed product). Noncrossed product division algebras were long conjectured to be fictional, until they were shown to exist by Amitsur in [1].

We say D is *indecomposable* if it does not contain a subalgebra that is also central over F , or equivalently if it is not an F -tensor product of two nontrivial F -division algebras. It is not hard to show that all division algebras of composite period are decomposable, and that all division algebras of equal prime-power period and index are indecomposable, but it is nontrivial to construct indecomposable division algebras of unequal prime-power period and index. The first examples appeared in [3] and in [31]. For additional discussion of either of these topics, see almost any survey treating open problems on division algebras, such as [5], [2], or [32].

We can use Theorem 5.4 to prove the existence of noncrossed product and indecomposable division algebras over the function field F of any p -adic curve $X_{\mathbb{Q}_p}$. Noncrossed products over $K(t)$ for K a local field were first constructed in [8], and then constructed more systematically over the function field of a smooth relative \mathbb{Z}_p -curve in [10]. Indecomposable division algebras of unequal period and index were also constructed in [10], over the same types of fields. Modulo gluing, the method we use below is the same as the one used in [10, Theorem 4.3, Corollary 4.8].

Theorem 6.1. *Let F/\mathbb{Q}_p be a finitely generated field extension of transcendence degree one. Let X/\mathbb{Z}_p be a regular relative curve with function field F , let C_i be a reduced irreducible component of the closed fiber, let $\ell \neq p$ be a prime, and let r and s be numbers that are maximal such that $\mu_{\ell^r} \subset \kappa(C_1)$ and $\mu_{\ell^s} \subset \kappa(C_1)(\mu_{\ell^{r+1}})$. Then there exist noncrossed product F -division algebras of period and index as low as ℓ^{s+1} if $r = 0$, and ℓ^{2r+1} if $r \neq 0$.*

Proof. We may assume (without changing r and s) that C has regular irreducible components, at most two of which meet at any closed point of X . The idea is to use the (known) existence of such algebras over the fields F_{C_i} , modify the construction to produce a class in $H^2(\mathcal{O}_{C,S}) \oplus \Gamma$, and then apply Theorem 4.6 and Theorem 5.4 to prove existence over F .

By [10, Theorem 4.7], if F admits a smooth X then there exist noncrossed product division algebras over F_C of period and index as low as ℓ^{s+1} if $r = 0$, and ℓ^{2r+1} if $r > 0$. The resulting Brauer class has the form $\alpha_C + (\pi) \cdot \theta_C \in H^2(F_C)$, where $\alpha_C \in H^2(\kappa(C))$ and $\theta_C \in H^1(\kappa(C), -1)$. A look at the construction, which proceeds exactly as in [6, Theorem 1], shows we may pre-assign values at any finite set of points of C . Thus we may thus produce a noncrossed product F_{C_1} -division algebra with class $\gamma_{C_1} = \alpha_{C_1} + (\pi) \cdot \theta_{C_1}$ of the desired period and index, and elements $\gamma_{C_i} = \alpha_{C_i} + (\pi) \cdot \theta_{C_i}$ for $i > 1$ that glue as in Lemma 4.2. Let

$$\gamma_C = (\alpha_{C_1} + (\pi) \cdot \theta_{C_1}, \dots, \alpha_{C_m} + (\pi) \cdot \theta_{C_m}) \in H^2(\mathcal{O}_{C,S}) \oplus (\pi) \cdot \Gamma \leq \bigoplus_i H^2(F_{C_i})$$

Then γ_C lifts to $\gamma = \lambda(\gamma_C) \in H^2(F)$ by Theorem 4.6. By Theorem 5.4, $\text{ind}(\gamma) = \text{ind}(\gamma_C) = \ell^{s+1}$ if $r = 0$, and ℓ^{2r+1} if $r \neq 0$. It is clear that the F -division algebra

D associated to γ is a noncrossed product, since any Galois maximal subfield of D over F could be constructed over F_{C_1} , contradicting the fact that $D \otimes_F F_{C_1}$ is a noncrossed product F_{C_1} -division algebra. ■

Theorem 6.2. *Let F/\mathbb{Q}_p be a finitely generated field extension of transcendence degree one, and let $\ell \neq p$ be a prime. Then there exist indecomposable F -division algebras of $(\text{period}, \text{index}) = (\ell^a, \ell^b)$, for any numbers a and b satisfying $1 \leq a \leq b \leq 2a - 1$.*

Proof. Let X , C , C_i , and \mathcal{S} be as in Theorem 6.1. The construction over F_{C_i} is exactly as in [10, Proposition 4.2] and [7], and we merely have to observe that we may assume all of the data in the constructed class $\gamma_{C_i} = \alpha_{C_i} + (\pi) \cdot \theta_{C_i} \in H^2(F_{C_i})$ is trivial at the singular points $\mathcal{S} \cap C_i$, so that by Lemma 4.2, we may construct a class $\gamma_C = \alpha_C + (\pi) \cdot \theta_C$ in $H^2(\text{O}_{C, \mathcal{S}}) \oplus (\pi) \cdot \Gamma \leq \bigoplus_i H^2(F_{C_i})$ whose i -th component is $\alpha_{C_i} + (\pi) \cdot \theta_{C_i}$. This class lifts to a class $\gamma = \lambda(\gamma_C)$ by Theorem 4.6, and $\text{ind}(\gamma) = \text{ind}(\gamma_C)$ by Theorem 5.4. Since the indexes are the same, the division algebra D associated to γ is indecomposable, since any decomposition would extend to $D_{C_i} = D \otimes_F F_{C_i}$, contradicting the construction of γ_{C_i} . ■

REFERENCES

- [1] S. Amitsur. On central division algebras. *Israel J. Math.*, 12:408–422, 1972.
- [2] S. Amitsur. Division algebras. A survey. In *Algebraists' homage: papers in ring theory and related topics (New Haven, Conn., 1981)*, volume 13 of *Contemp. Math.*, pages 3–26. Amer. Math. Soc., Providence, R.I., 1982.
- [3] S. Amitsur, L. Rowen, and J.-P. Tignol. Division algebras of degree 4 and 8 with involution. *Israel J. Math.*, 33:133–148, 1979.
- [4] M. F. Atiyah and I. G. Macdonald. *Introduction to Commutative Algebra*. Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.
- [5] A. Auel, E. Brussel, S. Garibaldi, and U. Vishne. Open problems on central simple algebras. *Transform. Groups*, 16(1):219–264, March 2011.
- [6] E. Brussel. Noncrossed products and nonabelian crossed products over $\mathbb{Q}(t)$ and $\mathbb{Q}((t))$. *Amer. J. Math.*, 117(2):377–393, 1995.
- [7] E. Brussel. Decomposability and embeddability of discretely Henselian division algebras. *Israel J. Math.*, 96(part A):141–183, 1996.
- [8] E. Brussel. Noncrossed products over $k_p(t)$. *Trans. Amer. Math. Soc.*, 353(5):2115–2129 (electronic), 2001.
- [9] E. Brussel. On Saltman's p -adic curves papers. In *Quadratic forms, linear algebraic groups, and cohomology*, volume 18 of *Dev. Math.*, pages 13–39. Springer, New York, 2010.
- [10] E. Brussel, K. McKinnie, and E. Tengan. Indecomposable and noncrossed product division algebras over function fields of smooth p -adic curves. *Adv. in Math.*, 226:4316–4337, 2011.
- [11] E. Brussel and E. Tengan. Division algebras of prime period $\ell \neq p$ over function fields of p -adic curves. [arxiv.org:1003.1906](https://arxiv.org/abs/1003.1906), 2010.
- [12] J.-L. Colliot-Thélène. Birational invariants, purity and the Gersten conjecture. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 1–64. Amer. Math. Soc., Providence, RI, 1995.
- [13] E. Freitag and R. Kiehl. *Étale cohomology and the Weil conjecture*, volume 13 of *Ergebnisse der Mathematik und ihrer Grenzgebiete (3) [Results in Mathematics and Related Areas (3)]*. Springer-Verlag, Berlin, 1988. Translated from the German by Betty S. Waterhouse and William C. Waterhouse, With an historical introduction by J. A. Dieudonné.

- [14] L. Fuchs. *Infinite Abelian Groups. Vol. I*. Pure and Applied Mathematics, Vol. 36. Academic Press, New York, 1970.
- [15] K. Fujiwara. A proof of the absolute purity conjecture (after Gabber). In *Algebraic geometry 2000, Azumino (Hotaka)*, volume 36 of *Adv. Stud. Pure Math.*, pages 153–183. Math. Soc. Japan, Tokyo, 2002.
- [16] S. Garibaldi, A. Merkurjev, and J.-P. Serre. *Cohomological invariants in Galois cohomology*, volume 28 of *University Lecture Series*. Amer. Math. Soc., 2003.
- [17] A. Grothendieck. Éléments de géométrie algébrique. I. Le langage des schémas. *Inst. Hautes Études Sci. Publ. Math.*, (4):228, 1960.
- [18] A. Grothendieck. Éléments de géométrie algébrique. II. Étude globale élémentaire de quelques classes de morphismes. *Inst. Hautes Études Sci. Publ. Math.*, (8):222, 1961.
- [19] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. II. *Inst. Hautes Études Sci. Publ. Math.*, (24):231, 1965.
- [20] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III. *Inst. Hautes Études Sci. Publ. Math.*, (28):255, 1966.
- [21] A. Grothendieck. Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas IV. *Inst. Hautes Études Sci. Publ. Math.*, (32):361, 1967.
- [22] A. Grothendieck. *Revêtements étales et groupe fondamental (SGA 1)*. Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 3. Société Mathématique de France, Paris, 2003. Séminaire de géométrie algébrique du Bois Marie 1960–61. [Algebraic Geometry Seminar of Bois Marie 1960–61], Directed by A. Grothendieck, With two papers by M. Raynaud, Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)].
- [23] A. Grothendieck and J. P. Murre. *The tame fundamental group of a formal neighbourhood of a divisor with normal crossings on a scheme*. Lecture Notes in Mathematics, Vol. 208. Springer-Verlag, Berlin, 1971.
- [24] T. Hanke. A twisted Laurent series ring that is a noncrossed product. *Israel J. Math.*, 150:199–203, 2005.
- [25] S. Lang. *Algebra*. Springer-Verlag, third edition, 2002.
- [26] Q. Liu. *Algebraic Geometry and Arithmetic Curves*, volume 6 of *Oxford Graduate Texts in Mathematics*. Oxford University Press, Oxford, 2002. Translated from the French by Reinie Ern , Oxford Science Publications.
- [27] H. Matsumura. *Commutative ring theory*, volume 8 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, Cambridge, second edition, 1989. Translated from the Japanese by M. Reid.
- [28] J. Milne. * tale Cohomology*. Princeton University Press, 1980.
- [29] W. Raskind. Abelian class field theory of arithmetic schemes. In *K-theory and algebraic geometry: connections with quadratic forms and division algebras (Santa Barbara, CA, 1992)*, volume 58 of *Proc. Sympos. Pure Math.*, pages 85–187. Amer. Math. Soc., Providence, RI, 1995.
- [30] S. Saito. Class field theory for curves over local fields. *J. Number Theory*, 21(1):44–80, 1985.
- [31] D. Saltman. Indecomposable division algebras. *Comm. Algebra*, 7(8):791–817, 1979.
- [32] D. Saltman. Finite dimensional division algebras. In *Azumaya algebras, actions, and modules*, volume 124 of *Contemp. Math.*, pages 203–214. Amer. Math. Soc., 1992.
- [33] D. Saltman. Division algebras over p -adic curves. *J. Ramanujan Math. Soc.*, 12:25–47, 1997. see also the erratum [34] and survey [9].
- [34] D. Saltman. Correction to division algebras over p -adic curves. *J. Ramanujan Math. Soc.*, 13:125–129, 1998.
- [35] D. Saltman. Cyclic algebras over p -adic curves. *J. Algebra*, 314:817–843, 2007.
- [36] V. Suresh. Bounding the symbol length in the Galois cohomology of function fields of p -adic curves. *Comment. Math. Helv.*, 85(2):337–346, 2010.

DEPARTMENT OF MATHEMATICS & COMPUTER SCIENCE, EMORY UNIVERSITY, ATLANTA, GA 30322, USA

E-mail address: brussel@mathcs.emory.edu

INSTITUTO DE CIÊNCIAS MATEMÁTICAS E DE COMPUTAÇÃO, UNIVERSIDADE DE SÃO PAULO, SÃO
CARLOS, SÃO PAULO, BRAZIL

E-mail address: `etengan@icmc.usp.br`